

# JAAM FORECAST sepsis②

Wada et al. *Journal of Intensive Care* (2023) 11:8  
<https://doi.org/10.1186/s40560-023-00656-5>

Journal of Intensive Care

RESEARCH

Open Access



Sepsis-related coagulopathy treatment based on the disseminated intravascular coagulation diagnostic criteria: a post-hoc analysis of a prospective multicenter observational study

Takeshi Wada<sup>1\*</sup>, Kazuma Yamakawa<sup>2</sup>, Daijiro Kabata<sup>3</sup>, Toshikazu Abe<sup>4,5</sup>, Seitaro Fujishima<sup>6</sup>, Shigeki Kushimoto<sup>7</sup>, Toshihiko Mayumi<sup>8</sup>, Hiroshi Ogura<sup>9</sup>, Daizoh Saitoh<sup>10</sup>, Atsushi Shiraishi<sup>11</sup>, Yasuhiro Otomo<sup>12</sup> and Satoshi Gando<sup>1,13</sup> on behalf of the JAAM FORECAST Group

## Target: DICでかつ重症

→JAAM基準+PTで評価できないか

- ◆ PTが高値でDICが重症になるほど死亡、多臓器不全↑  
(PT-INR1.5あたりまでが顕著)
- ◆ DIC重症かつPT-INR高くなるにつれ抗凝固療法の生存率改善効果が顕著
- ◆ その閾値は、**JAAM 5かつPT-INT 1.5**

